



The NOvA Experiment

Ralf Ehrlich

(for the NOvA collaboration)

University of Virginia

June 21st, 2012





Overview

- The name
 - N** NuMI = Neutrinos at the Main Injector
 - O** Off axis
 - v** ν_e
 - A** Appearance
- 150+ Collaborators in 25 Institutions from 5 Countries
- The NOvA experiment
 - long baseline neutrino experiment
 - Designed to look for $\nu_\mu \rightarrow \nu_e$ oscillations
 - consists of two detectors
 - The Near Detector located at Fermilab
 - The Far Detector located in Northern Minnesota
 - 810 km away from each other



The Neutrino Source


- 120 GeV protons are extracted from Fermilab's Main Injector and get fired onto a graphite target
- Results in the creation of charged particles – in particular pions
- A magnetic horn focuses the charged pions towards the Soudan Mine (MINOS Far Detector)
- The pions decay into (anti-) muons and (anti-) muon-neutrinos traveling in approximately the same direction
- Reversal of the horn current is used to switch between neutrino and anti-neutrino mode



- Currently shut down for an upgrade until April 2013
 - will deliver $4.9 \cdot 10^{13}$ protons every 1.3 s for 10 μ s
 - will have a beam power of 700 kW (was 380 kW before shot down)


Detector Design

- The detectors are built of plastic cells
 - 4.0 cm x 6.7 cm x 15 m (Near Detector cells are shorter)
 - 32 cells form one module
 - modules arranged in alternating horizontal and vertical planes to enable a 3D reconstruction
- Cells are filled with liquid scintillator
- Charged particles due to neutrino interactions causes the scintillator to emit light
- This light gets captured by optical fibers
- Optical fibers transmit the light to Avalanche Photo Diodes (APD)
 - 85% quantum efficiency
 - Operated at 425 V
 - Cooled to $-15\text{ }^{\circ}\text{C}$ by thermoelectric coolers
- The APD output is digitized at the Front End Boards in the electronics box at the end of manifold covers.




The Prototype Near Detector

- Prototype detector
 - Built to get some experience before building the “real” Near Detector and Far Detector
 - Has been invaluable for developing of reconstruction algorithms and prototyping the equipment
- Located on the surface at Fermilab 1km away from the target
- Has a “Muon Catcher” made of steel to contain muons from CC interactions
- 16,000 cells, 222 tons
- Status
 - Finished in spring 2011
 - First events recorded in fall 2010





The Prototype Near Detector




The Near Detector

- The “real” Near Detector
- Located underground at Fermilab 1 km away from the target
- Identical design as the Far Detector – except that it is smaller
- Has a “Muon Catcher” made of steel to contain muon tracks from CC interactions
- 20,000 cells, 266 tons
- Status:
 - Construction starts in 2013
 - Construction finished in 2014
- Will be used to measure the ν_μ flux emitted at the source of the neutrino beam
- Will be used to study background events:
 - ν_e contaminations of the neutrino beam
 - neutral current ν_μ interactions




The Far Detector

- Located in Northern Minnesota 810 km away from the neutrino source
- 356,000 cells, 14,000 tons, 15x15x63 m³
- Status:
 - Building dedication in April 2012
 - Block installation will start in summer 2012
 - Construction will be finished in 2014
- Designed to detect the ν_e after they oscillated from ν_μ
- Sits near the first expected $\nu_\mu \rightarrow \nu_e$ oscillation peak



The Far Detector




The Far Detector



Off Axis Location


- Near and Far Detectors are located 14 mrad away from the beam axis
 - At this angle, the neutrino beam has a narrow energy distribution with a peak around 2 GeV
 - Reduces the background
- The transition probability for $\nu_\mu \rightarrow \nu_e$ is expected to be close to its maximum
 - at this energy (2 GeV) and
 - at this location (810 km away from the neutrino source)
- The neutrino beam goes through the Earth so that we can make use of the matter effects




Neutrino Detection

Typical (simulated) events


- Long μ tracks
- Short proton track at the vertex may be visible



- Short electron tracks with EM cascades along the track
- Short proton track at the vertex may be visible



- $\pi^0 \rightarrow \gamma\gamma$
- Identified by gap b/w beginning and end of the γ tracks



Neutrino Mixing Parameters

Global status of neutrino oscillation parameters after recent reactor measurements

from
arXiv:1205.4018
v2 [hep-ph]
21 May 2012

We don't know the sign, since we don't know the mass hierarchy

parameter	best fit $\pm 1\sigma$	2σ	3σ
$\Delta m_{21}^2 [10^{-5} \text{eV}^2]$	7.62 ± 0.19	7.27–8.01	7.12–8.20
$\Delta m_{31}^2 [10^{-3} \text{eV}^2]$	$2.53^{+0.08}_{-0.10}$ $-(2.40^{+0.10}_{-0.07})$	2.34 – 2.69 $-(2.25 - 2.59)$	2.26 – 2.77 $-(2.15 - 2.68)$
$\sin^2 \theta_{12}$	$0.320^{+0.015}_{-0.017}$	0.29–0.35	0.27–0.37
$\sin^2 \theta_{23}$	$0.49^{+0.08}_{-0.05}$ $0.53^{+0.05}_{-0.07}$	0.41–0.62 0.42–0.62	0.39–0.64
$\sin^2 \theta_{13}$	$0.026^{+0.003}_{-0.004}$ $0.027^{+0.003}_{-0.004}$	0.019–0.033 0.020–0.034	0.015–0.036 0.016–0.037
δ	$(0.83^{+0.54}_{-0.64}) \pi$ $0.07\pi^a$	$0 - 2\pi$	$0 - 2\pi$

Is $\sin^2(2\theta_{23}) = 1.0$,
i.e. $\theta_{23} = 45^\circ$?


$\theta_{13} \approx 9^\circ$

In other words:
unknown

TABLE I: Neutrino oscillation parameters summary. For Δm_{31}^2 , $\sin^2 \theta_{23}$, $\sin^2 \theta_{13}$, and δ the upper (lower) row corresponds to normal (inverted) neutrino mass hierarchy.

Neutrino Mixing Parameters

- We measure the appearance of ν_e and $\bar{\nu}_e$ i.e. $P(\nu_\mu \rightarrow \nu_e)$ and $P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$ which are functions of
 - mixing angles $\theta_{13}, \theta_{23}, \theta_{12}$
 - the CP violating phase δ
 - the mass hierarchy, i.e. normal or inverted
 - mass squared differences $\Delta m_{31}^2, \Delta m_{21}^2, \Delta m_{32}^2$
 - neutrino energy (2 GeV)
 - oscillation distance (810 km)
 - matter effects




- Approximated probability function

$$\begin{aligned}
 - P_{vac} \left(\begin{array}{l} \nu_\mu \rightarrow \nu_e \\ \bar{\nu}_\mu \rightarrow \bar{\nu}_e \end{array} \right) &\approx \sin^2(\theta_{23}) \sin^2(2\theta_{13}) \sin^2(\Delta_{31}) \\
 &+ \cos^2(\theta_{13}) \cos^2(\theta_{23}) \sin^2(2\theta_{12}) \sin^2(\Delta_{21}) \\
 &+ J \sin(\Delta_{21}) \sin(\Delta_{31}) [\cos(\Delta_{32}) \cos(\delta) \mp \sin(\Delta_{32}) \sin(\delta)] \\
 - \text{with } \Delta_{ij} &= 1.27 \frac{\Delta m_{ij}^2 L}{E} \text{ and } \Delta m_{ij} \text{ in eV}^2, L \text{ in km, and } E \text{ in GeV} \\
 - \text{and } J &= \cos(\theta_{13}) \sin(\theta_{12}) \sin(\theta_{13}) \sin(\theta_{23}) \\
 - \text{matter effect } P_{matter} \left(\begin{array}{l} \nu_\mu \rightarrow \nu_e \\ \bar{\nu}_\mu \rightarrow \bar{\nu}_e \end{array} \right) &\approx \left(1 \pm \frac{E}{6\text{Gev}} \right) P_{vac} \left(\begin{array}{l} \nu_\mu \rightarrow \nu_e \\ \bar{\nu}_\mu \rightarrow \bar{\nu}_e \end{array} \right)
 \end{aligned}$$


↖ Reversed for inverted mass hierarchy

Neutrino Mixing Parameters

- NOvA measures the appearance probabilities for ν_e and $\bar{\nu}_e$
- The plot show how one may be able to extract θ_{13} , the mass hierarchy and the CP violating phase δ
 - This becomes more difficult for smaller values of θ_{13} due to overlaps of both ellipses
 - Fortunately, θ_{13} may be large according to the latest results
- The plot will look differently if $\sin(2\theta_{23}) < 1$



Neutrino Mixing Parameters





normal
hierarchy

inverted
hierarchy

NOvA


The mass hierarchy can be resolved with better than 95% C.L. for values to the right of the curves.



normal
hierarchy

inverted
hierarchy

**NOvA +
T2K**



Neutrino Mixing Parameters

- θ_{23} and Δm_{32}^2 can be determined by looking at the ν_μ survival probability $P(\nu_\mu \rightarrow \nu_\mu) \approx 1 - \sin^2(2\theta_{23})\sin^2(\Delta_{32})$

with $\Delta_{ij} = 1.27 \frac{\Delta m_{ij}^2 L}{E}$ and Δm_{ij} in eV^2 , L in km, and E in GeV


- This approximation assumes that the ν_μ disappearance is mainly caused by $\nu_\mu \rightarrow \nu_\tau$ (see arXiv:0711.0769v1 [hep-ex])
- P depends on θ_{23} , Δm_{32}^2 , L , E
- P can be determined as a function of E

by comparing the ν_μ energy spectrum at the ND and FD

- Since L is known, this leaves the two unknown variables θ_{23} and Δm_{32}^2 which can be fitted to P to obtain their values
 - However, we only get $\sin^2(2\theta_{23})$ and cannot distinguish whether $\theta_{23} > 45^\circ$ or $\theta_{23} < 45^\circ$
- Requires an excellent energy resolution for muons (as a product of the CC ν_μ reaction)

Neutrino Mixing Parameters

- If $\sin^2(2\theta_{23}) < 1 \rightarrow \theta_{23} \neq 45^\circ$
 - We need to determine whether θ_{23} is less or greater than 45°
 - The $P(\nu_e) - P(\bar{\nu}_e)$ map will look differently
 - Reactor experiments (e.g. Daya Bay) measure $P(\bar{\nu}_e \rightarrow \bar{\nu}_e)$ which depends on $\sin^2(2\theta_{13})$
 - unlike NOvA which depends on $\sin^2(\theta_{23})\sin^2(2\theta_{13})$
 - Comparison can be used to determine whether θ_{23} is less or greater than 45°



Other Physics Opportunities

- Magnetic monopoles
 - can be identified as highly ionizing and/or very slow moving particles going through the entire detector
- Supernova neutrinos
 - the detector will be subjected to a larger amount of neutrinos coinciding with a supernova
- Solar WIMPs
 - would result in high energetic neutrino events coming from the direction of the sun

Summary

- Many physics goals can be reached
 - Measuring the neutrino mixing angles θ_{13} , θ_{23}
 - Measuring the mass squared difference Δm_{32}^2
 - Determining the CP violating complex phase δ
 - Determining the mass hierarchy
- Recent θ_{13} results show that it is large, which helps us to reach our physics goals
- The Prototype Near Detector at the surface
 - Has already been running successfully
 - Has provided us with valuable information about the detector construction
- Far Detector construction will begin this summer
 - First events can be recorded as soon as the first block is installed
 - First oscillation events can be observed as soon as the neutrino beam gets turned on in April 2013




Thank You

Backup Slides

Neutrino Mixing Parameters


- Example:
 - Assume we measure
$$P(\nu_\mu \rightarrow \nu_e) = 0.025$$
$$P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) = 0.060$$
 - 3 years in neutrino mode and 3 years in anti-neutrino mode



Neutrino Mixing Parameters

NOVA hierarchy resolution, 3+3 yr ($\nu+\bar{\nu}$)

$\sin^2(2\theta_{13})=0.095$, $\sin^2(2\theta_{23})=1.00$



Neutrino Mixing Parameters

